



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2011

Vitamin D: role in pregnancy and early childhood

Bischoff-Ferrari, H A

Abstract: Several studies in pregnant women and early childhood suggest that vitamin D deficiency (serum 25-hydroxyvitamin D levels <50 nmol/l) is common in both population groups. Recent recommendations have therefore reviewed the literature regarding the role of vitamin D in pregnant women and in early childhood. The Institute of Medicine, in their most recent assessment in 2010, recommended 600 IU per day in pregnant and lactating women. In 2011, the US Endocrine Task Force on Vitamin D commented that 600 IU per day may not be sufficient to correct vitamin D deficiency in pregnant and lactating women. Their recommendation was 1,500-2,000 IU vitamin D per day in pregnant and lactating women with vitamin D deficiency. For infants, the recommendation from both societies is consistently 400 IU vitamin D per day, and also in children both societies recommend 600 IU vitamin D per day. This review will summarize the scientific basis that led to the most recent recommendations.

DOI: <https://doi.org/10.1159/000332069>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-59160>

Journal Article

Published Version

Originally published at:

Bischoff-Ferrari, H A (2011). Vitamin D: role in pregnancy and early childhood. *Annals of Nutrition Metabolism*, 59(1):17-21.

DOI: <https://doi.org/10.1159/000332069>

Vitamin D – Role in Pregnancy and Early Childhood

Heike A. Bischoff-Ferrari

Centre on Aging and Mobility, University of Zurich and City Hospital Waid, University Hospital, Zurich, Switzerland

Key Words

Vitamin D • Pregnancy • Supplement • Recommendation • Infancy • Recommended intake

Abstract

Several studies in pregnant women and early childhood suggest that vitamin D deficiency (serum 25-hydroxyvitamin D levels <50 nmol/l) is common in both population groups. Recent recommendations have therefore reviewed the literature regarding the role of vitamin D in pregnant women and in early childhood. The Institute of Medicine, in their most recent assessment in 2010, recommended 600 IU per day in pregnant and lactating women. In 2011, the US Endocrine Task Force on Vitamin D commented that 600 IU per day may not be sufficient to correct vitamin D deficiency in pregnant and lactating women. Their recommendation was 1,500–2,000 IU vitamin D per day in pregnant and lactating women with vitamin D deficiency. For infants, the recommendation from both societies is consistently 400 IU vitamin D per day, and also in children both societies recommend 600 IU vitamin D per day. This review will summarize the scientific basis that led to the most recent recommendations.

Copyright © 2011 S. Karger AG, Basel

Threshold Definition for Vitamin D Deficiency and Adequacy

As a general guideline, vitamin D deficiency has been defined as a 25-hydroxyvitamin D level [25(OH)D] of less than 50 nmol/l (<20 ng/ml) in all age groups [1]. At levels below 50 nmol/l, bone resorption and increased PTH levels have been documented [2]. Levels lower than 25 nmol/l (<10 ng/ml) are considered to indicate severe deficiency and can induce adverse effects such as rickets in infants and osteomalacia in adults [2]. Vitamin D adequacy needs to be established in children and pregnant women; however, observational data in younger (age 19–49 years), middle-aged (age 50–64 years), and senior adults (age ≥ 65 years) suggest that a level of no less than 75 nmol/l is needed for optimal hip bone density [3], and this threshold is consistent with fracture prevention data from double-blind randomized trials in senior adults [4].

Prevalence of Vitamin D Deficiency in Pregnant Women

Surveys of pregnant women in Greece [5], Belgium [6], and the Netherlands [7] suggest that 10–30% of pregnant women have 25(OH)D levels below 30 nmol/l.

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2011 S. Karger AG, Basel
0250–6807/11/0591–0017\$38.00/0

Accessible online at:
www.karger.com/anm

Prof. Heike A. Bischoff-Ferrari, MD, DrPH
Centre on Aging and Mobility, University of Zurich and City Hospital Waid
University Hospital, Gloriastrasse 25
CH–8091 Zurich (Switzerland)
E-Mail HeikeABischoff@aol.com

While the threshold of 50 nmol/l was not assessed in the available European studies, the Belgium study documented that 86–88 % of pregnant women did not reach 25(OH)D levels of at least 75 nmol/l [6]. Furthermore, in the Dutch study, 59–84% of pregnant women of non-Western ethnicity had 25(OH)D levels below 30 nmol/l [7]. This high prevalence among pregnant women of non-Western ethnicity may be explained either by a darker skin tone (natural sun protection) or the avoidance of direct sunshine exposure for cultural reasons [8]. The Greek study also measured 25(OH)D levels in newborns and found that their levels were strongly correlated with those of their mothers (correlation coefficient = 0.64), suggesting that the mother's vitamin D status is important for the vitamin D status of the newborn [5]. The high prevalence of vitamin D deficiency among pregnant women led the US Endocrine Task Force on Vitamin D to recommend 25(OH)D assessment in all pregnant women [9].

Prevalence of Vitamin D Deficiency in Early Childhood

During lactation, the newborn largely depends on nutritional sources of vitamin D through breast milk, fortified newborn nutrition, or supplements. To satisfy the requirements of an infant who is only fed breast milk, the mother requires 2,000–4,000 IU per day to transfer enough vitamin D into her milk [10]. Alternatively, as recommended in most countries, newborns should receive a supplement of 400–500 IU vitamin D per day. In a larger survey of German children age 1–17 years, median 25(OH)D levels were 44 nmol/l among native children and 35 nmol/l among immigrant children. The levels of boys and girls were similar; however, there was some decline with age, with the highest median levels measured in the first year of life, likely due to the German supplementation recommendation of 500 IU per day in infants (KiGGS [11]). The Optiford study of 4 Northern European countries (Denmark, Finland, Poland, and Ireland) included 199 children with a mean age of 12.5 years who had serum 25(OH)D levels below 25 nmol/l in 30–50% of cases and serum levels below 48 nmol/l in over 90% of cases [12]. The same study also assessed 25(OH)D levels in seniors and suggested that vitamin D deficiency was more prevalent in children [12].

Why Is It Important to Avoid Vitamin D Deficiency in Pregnant Women? – Relevance for Mother and Child

Relevance for Mothers

Vitamin D deficiency during pregnancy has been associated with an increased risk of preeclampsia [13], gestational diabetes [14], and caesarian section [5], but data from intervention studies are missing.

Relevance for Children

Observational data suggest that increased vitamin D exposure of the mother during pregnancy correlates with better bone health in the offspring. In a longitudinal study of 198 children in the UK, reduced 25(OH)D levels in their mothers during late pregnancy was associated with reduced whole-body ($r = 0.21$, $p = 0.009$) and lumbar spine ($r = 0.17$, $p = 0.03$) bone mineral content in children at age 9 years [15]. This may in part be explained by some data from observational studies [16–19] and small clinical trials [20, 21] suggesting that a higher vitamin D status among pregnant women correlates positively with the birth weight of their offspring, although some observational studies showed a neutral association [22–26]. In a small double-blind randomized controlled trial among 126 Asian women living in London, supplementation with 1,000 IU vitamin D₂ per day compared to placebo in the last trimester of pregnancy led to a greater weight gain in the mothers and reduced the risk of small-for-gestational-age babies (15 vs. 29%) [20]. These findings were supported by another study in Indian women and their offspring comparing a bolus supplementation with 600,000 IU vitamin D in gestational months 7 and 8 and 1,200 IU vitamin D per day in the third trimester to placebo, resulting in significantly higher birth weights with supplementation (bolus vitamin D = 3,180 g; daily vitamin D = 2,890 g; placebo = 2,730 g) [21]. No difference in birth weight was found in a smaller open-design trial comparing two treatment regimens with vitamin D to placebo (200,000 IU once during pregnancy or 1,000 IU per day during the last 3 months of pregnancy) [27]. The largest double-blind randomized controlled trial among 494 Caucasian, Hispanic, and African American pregnant women living in South Carolina by Hollis et al. [28] compared 400 IU to 2,000 IU and to 4,000 IU vitamin D per day. The mean 25(OH)D levels at baseline were about 60 nmol/l, and there was a significant increase in the mean 25(OH)D levels with treatment in all groups at 1 month prior to delivery (400 IU = 81.2 nmol/l; 2,000 IU = 102.6 nmol/l; 4,000 IU = 114.2 nmol/l). With respect to clinical end

points, no adverse events occurred with any of the tested doses, and small but nonsignificant differences in birth weight (400 IU = 3,222 g; 2,000 IU = 3,360 g; 4,000 IU = 3,285 g) and gestational age at birth (400 IU = 38.6 weeks; 2,000 IU = 38.8 weeks; 4,000 IU = 39.1 week) were documented [28].

There is also observational data suggesting that children exposed to more vitamin D during pregnancy may have a lower risk of developing asthma in early life as suggested by two longitudinal studies [29, 30]. Litonjua et al. [31] estimated from two birth cohort studies that the population attributable risk of asthma incidence caused by vitamin D deficiency in pregnancy is about 40% of all cases in children 3–5 years old. In contrast, one study suggested an increased risk of eczema on examination at 9 months (OR 3.26; 95% CI 1.15–9.29) and of asthma at age 9 years (OR 5.40; 95% CI 1.09–26.65) observed in a smaller cohort study among children of mothers who had 25(OH)D levels ≥ 75 nmol/l compared to children whose mothers had a concentration < 30 nmol/l during pregnancy [24]. The study has been criticized for its large loss to follow-up (62%) [24].

Why Is It Important to Avoid Vitamin D Deficiency in Early Childhood?

Bone Health

Rickets is a result of severe vitamin D deficiency and has a significant impact on growth and bone development [32, 33]. While supplementation with 100, 200, or 400 IU per day of vitamin D resulted in the prevention of rickets in one study [34], vitamin D intakes between 340 and 600 IU per day have been reported to have the maximum effect on the linear growth of infants [35]. Furthermore, adding to the concept that vitamin D supplementation in children may not only prevent rickets but also contribute to bone growth and the achievement of an optimal peak bone mass, a 2011 meta-analysis of 4 double-blind randomized controlled trials of vitamin D supplementation in the range of 132–2,000 IU vitamin D per day compared to controls (placebo or lower dose) among a total of 639 children suggested a trend toward a small effect on lumbar spine bone mineral density in all treated children (standardized mean difference 0.15; 95% CI –0.01 to 0.31; $p = 0.07$). There was also a significant benefit of vitamin D supplementation in children with serum 25(OH)D levels below 35 nmol/l [36], with effects on the total body bone mineral content and lumbar spine bone mineral density equivalent to a 2.6 and

1.7% percentage point greater change from baseline in the supplemented group [36]. Notably, based on one randomized trial among Lebanese girls age 10–17 years included in this most recent meta-analysis, hip bone density increased more with 14,000 IU vitamin D per week (2,000 IU per day) compared to 1,400 IU per week (200 IU per day) at the 12-month follow-up without any report of toxicity and irrespective of baseline 25(OH)D levels [37].

Nonskeletal End Points

In the randomized trial among Lebanese girls comparing 14,000 IU vitamin D per week (2,000 IU per day) with 1,400 IU per week (200 IU per day), there was also a benefit on muscle mass measured by DEXA at the 12-month follow-up [37]. Regarding other nonskeletal benefits of vitamin D in children, observational data support an inverse relationship between a higher 25(OH)D status or vitamin D supplementation and a lower incidence of diabetes [38, 39], atopic allergies [40, 41], and upper respiratory infections [42]. However, evidence from clinical trials in children is very limited. One Finnish birth cohort study suggested that, in infants who received $\geq 2,000$ IU per day of vitamin D compared to no supplementation during the first year of life, there was an 86% reduction in the risk of developing type 1 diabetes in the following 31 years [43]. A recent meta-analysis of four case-control studies of about 6,500 infants found that supplemental vitamin D reduces the risk of type 1 diabetes [OR 0.71 (0.60–0.84)], with higher cumulative doses conferring a greater benefit [44]. Regarding infections, one double-blind randomized controlled clinical trial in 334 Japanese children who received 1,200 IU vitamin D per day compared to placebo documented a 42% reduction in influenza A [45]. A secondary analysis documented an 83% reduction in asthma attacks among children with asthma enrolled in the same trial [45].

Recommendations

The Institute of Medicine, in their most recent assessment in 2010, recommended 600 IU per day in pregnant and lactating women [1]. In 2011, the US Endocrine Task Force on Vitamin D commented that 600 IU per day may not be sufficient to correct vitamin D deficiency in pregnant and lactating women [9]. Their recommendation is

1,500–2,000 IU vitamin D per day in pregnant and lactating women with vitamin D deficiency [9]. For infants, the recommendation of both societies is consistently 400 IU vitamin D per day, and also in children both societies recommend 600 IU vitamin D per day.

Disclosure Statement

The author has nothing to disclose.

References

- 1 IOM: Dietary reference ranges for calcium and vitamin D. 2010. <http://www.iom.edu/Reports/2010/Dietary-Reference-Intakes-for-Calcium-and-Vitamin-D/Report-Brief.aspx>.
- 2 Holick MF: Vitamin D deficiency. *N Engl J Med* 2007;357:266–281.
- 3 Bischoff-Ferrari HA, Dietrich T, Orav EJ, Dawson-Hughes B: Positive association between 25-hydroxy vitamin D levels and bone mineral density: a population-based study of younger and older adults. *Am J Med* 2004;116:634–639.
- 4 Bischoff-Ferrari HA, Willett WC, Wong JB, Stuck AE, Staehelin HB, Orav EJ, et al: Prevention of nonvertebral fractures with oral vitamin D and dose dependency: a meta-analysis of randomized controlled trials. *Arch Intern Med* 2009;169:551–561.
- 5 Merewood A, Mehta SD, Chen TC, Bauchner H, Holick MF: Association between vitamin D deficiency and primary cesarean section. *J Clin Endocrinol Metab* 2009;94:940–945.
- 6 Cavalier E, Delanaye P, Morreale A, Carlisi A, Mourad I, Chapelle JP, et al: Vitamin D deficiency in recently pregnant women (in French). *Rev Med Liege* 2008;63:87–91.
- 7 van der Meer IM, Karamali NS, Boeke AJ, Lips P, Middelkoop BJ, Verhoeven I, et al: High prevalence of vitamin D deficiency in pregnant non-Western women in The Hague, Netherlands. *Am J Clin Nutr* 2006;84:350–353; quiz 468–469.
- 8 Holick MF: Environmental factors that influence the cutaneous production of vitamin D. *Am J Clin Nutr* 1995;61(suppl):638S–645S.
- 9 Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al: Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96:1911–1930.
- 10 Hollis BW, Wagner CL: Vitamin D requirements during lactation: high-dose maternal supplementation as therapy to prevent hypovitaminosis D for both the mother and the nursing infant. *Am J Clin Nutr* 2004;80(suppl):1752S–1758S.
- 11 Hintzpetter B, Scheidt-Nave C, Muller MJ, Schenk L, Mensink GB: Higher prevalence of vitamin D deficiency is associated with immigrant background among children and adolescents in Germany. *J Nutr* 2008;138(8):1482–90.
- 12 Tylavsky FA, Cheng S, Lyytikäinen A, Viljakainen H, Lamberg-Allardt C: Strategies to improve vitamin D status in northern European children: exploring the merits of vitamin D fortification and supplementation. *J Nutr* 2006;136:1130–1134.
- 13 Bodnar LM, Catov JM, Simhan HN, Holick MF, Powers RW, Roberts JM: Maternal vitamin D deficiency increases the risk of pre-eclampsia. *J Clin Endocrinol Metab* 2007;92:3517–3522.
- 14 Zhang C, Qiu C, Hu FB, David RM, van Dam RM, Bralley A, et al: Maternal plasma 25-hydroxyvitamin D concentrations and the risk for gestational diabetes mellitus. *PLoS One* 2008;3:e3753.
- 15 Javaid MK, Crozier SR, Harvey NC, Gale CR, Dennison EM, Boucher BJ, et al: Maternal vitamin D status during pregnancy and childhood bone mass at age 9 years: a longitudinal study. *Lancet* 2006;367:36–43.
- 16 Leffelaar ER, Vrijkotte TG, van Eijsden M: Maternal early pregnancy vitamin D status in relation to fetal and neonatal growth: results of the multi-ethnic Amsterdam Born Children and their Development cohort. *Br J Nutr* 2010;104:1–10.
- 17 Bodnar LM, Catov JM, Zmuda JM, Cooper ME, Parrott MS, Roberts JM, et al: Maternal serum 25-hydroxyvitamin D concentrations are associated with small-for-gestational age births in white women. *J Nutr* 2010;140:999–1006.
- 18 Scholl TO, Chen X: Vitamin D intake during pregnancy: association with maternal characteristics and infant birth weight. *Early Hum Dev* 2009;85:231–234.
- 19 Sabour H, Hossein-Nezhad A, Maghbooli Z, Madani F, Mir E, Larijani B: Relationship between pregnancy outcomes and maternal vitamin D and calcium intake: a cross-sectional study. *Gynecol Endocrinol* 2006;22:585–589.
- 20 Brooke OG, Brown IR, Bone CD, Carter ND, Cleeve HJ, Maxwell JD, et al: Vitamin D supplements in pregnant Asian women: effects on calcium status and fetal growth. *Br Med J* 1980;280:751–754.
- 21 Marya RK, Rathee S, Lata V, Mudgil S: Effects of vitamin D supplementation in pregnancy. *Gynecol Obstet Invest* 1981;12:155–161.
- 22 Brunvand L, Quigstad E, Urdal P, Haug E: Vitamin D deficiency and fetal growth. *Early Hum Dev* 1996;45:27–33.
- 23 Farrant HJ, Krishnaveni GV, Hill JC, Boucher BJ, Fisher DJ, Noonan K, et al: Vitamin D insufficiency is common in Indian mothers but is not associated with gestational diabetes or variation in newborn size. *Eur J Clin Nutr* 2009;63:646–652.
- 24 Gale CR, Robinson SM, Harvey NC, Javaid MK, Jiang B, Martyn CN, et al: Maternal vitamin D status during pregnancy and child outcomes. *Eur J Clin Nutr* 2008;62:68–77.
- 25 Morley R, Carlin JB, Pasco JA, Wark JD: Maternal 25-hydroxyvitamin D and parathyroid hormone concentrations and offspring birth size. *J Clin Endocrinol Metab* 2006;91:906–912.
- 26 Prentice A, Jarjou LM, Goldberg GR, Bennett J, Cole TJ, Schoenmakers I: Maternal plasma 25-hydroxyvitamin D concentration and birthweight, growth and bone mineral accretion of Gambian infants. *Acta Paediatr* 2009;98:1360–1362.
- 27 Mallet E, Gugi B, Brunelle P, Henocq A, Basuyau JP, Lemeur H: Vitamin D supplementation in pregnancy: a controlled trial of two methods. *Obstet Gynecol* 1986;68:300–304.
- 28 Hollis BW, Johnson D, Hulsey TC, Ebeling M, Wagner CL: Vitamin D supplementation during pregnancy: Double blind, randomized clinical trial of safety and effectiveness. *J Bone Miner Res* 2011, DOI: [10.1002/jbmr.463](https://doi.org/10.1002/jbmr.463).
- 29 Camargo CA Jr, Rifas-Shiman SL, Litonjua AA, Rich-Edwards JW, Weiss ST, Gold DR, et al: Maternal intake of vitamin D during pregnancy and risk of recurrent wheeze in children at 3 y of age. *Am J Clin Nutr* 2007;85:788–795.
- 30 Devereux G, Litonjua AA, Turner SW, Craig LC, McNeill G, Martindale S, et al: Maternal vitamin D intake during pregnancy and early childhood wheezing. *Am J Clin Nutr* 2007;85:853–859.
- 31 Litonjua AA, Weiss ST: Is vitamin D deficiency to blame for the asthma epidemic? *J Allergy Clin Immunol* 2007;120:1031–1035.
- 32 Holick MF, Chen TC: Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr* 2008;87:1080S–1086S.
- 33 Prentice A, Goldberg GR, Schoenmakers I: Vitamin D across the lifecycle: physiology and biomarkers. *Am J Clin Nutr* 2008;88:500S–506S.

- 34 Specker BL, Ho ML, Oestreich A, Yin TA, Shui QM, Chen XC, et al: Prospective study of vitamin D supplementation and rickets in China. *J Pediatr* 1992;120:733–739.
- 35 Feliciano ES, Ho ML, Specker BL, Falciglia G, Shui QM, Yin TA, et al: Seasonal and geographical variations in the growth rate of infants in China receiving increasing dosages of vitamin D supplements. *J Trop Pediatr* 1994;40:162–165.
- 36 Winzenberg T, Powell S, Shaw KA, Jones G: Effects of vitamin D supplementation on bone density in healthy children: systematic review and meta-analysis. *BMJ* 2011; 342:c7254.
- 37 El-Hajj Fuleihan G, Nabulsi M, Tamim H, Maalouf J, Salamoun M, Khalife H, et al: Effect of vitamin D replacement on musculoskeletal parameters in school children: a randomized controlled trial. *J Clin Endocrinol Metab* 2006;91:405–412.
- 38 Erkkola M, Kaila M, Nwaru BI, Kronberg-Kippila C, Ahonen S, Nevalainen J, et al: Maternal vitamin D intake during pregnancy is inversely associated with asthma and allergic rhinitis in 5-year-old children. *Clin Exp Allergy* 2009;39:875–882.
- 39 Holick MF: Diabetes and the vitamin D connection. *Curr Diab Rep* 2008;8:393–398.
- 40 Back O, Blomquist HK, Hernell O, Stenberg B: Does vitamin D intake during infancy promote the development of atopic allergy? *Acta Derm Venereol* 2009;89:28–32.
- 41 Brehm JM, Celedon JC, Soto-Quiros ME, Avila L, Hunninghake GM, Forno E, et al: Serum vitamin D levels and markers of severity of childhood asthma in Costa Rica. *Am J Respir Crit Care Med* 2009;179:765–771.
- 42 Ginde AA, Mansbach JM, Camargo CA Jr: Vitamin D, respiratory infections, and asthma. *Curr Allergy Asthma Rep* 2009;9:81–87.
- 43 Hypponen E, Laara E, Reunanen A, Jarvelin MR, Virtanen SM: Intake of vitamin D and risk of type 1 diabetes: a birth-cohort study. *Lancet* 2001;358:1500–1503.
- 44 Zipitis CS, Akobeng AK: Vitamin D supplementation in early childhood and risk of type 1 diabetes: a systematic review and meta-analysis. *Arch Dis Child* 2008;93: 512–517.
- 45 Urashima M, Segawa T, Okazaki M, Kurihara M, Wada Y, Ida H: Randomized trial of vitamin D supplementation to prevent seasonal influenza A in schoolchildren. *Am J Clin Nutr* 2010;91:1255–1260.